

A Symmetry-Based Method for Constructing Nonlocally Related PDE Systems

George W. Bluman^{*} and Zhengzheng Yang[†]

*Department of Mathematics, University of British Columbia,
Vancouver, British Columbia V6T 1Z2, Canada*

Abstract: Nonlocally related partial differential equation (PDE) systems are useful in the analysis of a given PDE system. It is known that each local conservation law of a given PDE system systematically yields a nonlocally related system. In this paper, a new and complementary method for constructing nonlocally related systems is introduced. In particular, it is shown that each point symmetry of a given PDE system systematically yields a nonlocally related system. Examples include applications to nonlinear diffusion equations, nonlinear wave equations and nonlinear reaction-diffusion equations. As a consequence, previously unknown nonlocal symmetries are exhibited for two examples of nonlinear wave equations. Moreover, since the considered nonlinear reaction-diffusion equations have no local conservation laws, previous methods do not yield nonlocally related systems for such equations.

1. Introduction

An equivalent nonlocally related partial differential equation (PDE) system can play an important role in the analysis of a given PDE system. Each solution of such a nonlocally related PDE system yields a solution of the given PDE system and, conversely, each solution of the given PDE system yields a solution of the nonlocally related system. These corresponding solutions are obtained through connection formulas. More important, the relationship between the solutions is not one-to-one. Hence, for a given PDE system, one could be more successful when using a standard method of analysis, especially a coordinate independent method, to a nonlocally related PDE system. For instance, through a nonlocally related PDE system, one can systematically find nonlocal symmetries and nonlocal conservation laws of a given PDE system. It turns out that such nonlocal symmetries and nonlocal conservation laws arise as local symmetries and local conservation laws of the nonlocally related system. Thus any method depending on local symmetry analysis is valid for nonlocal related PDE systems. For example, one can construct an explicit non-invertible mapping that maps a given nonlinear PDE system to a linear PDE system if a nonlocally related system can be linearized. Consequently, it is of great interest to systematically find such nonlocally related systems for a given PDE system. A systematic procedure for finding nonlocally related PDE systems is presented in [1] and references therein. Here one constructs a tree of nonlocally related systems that consists of potential systems and subsystems. The potential systems arise naturally from local conservation laws. However, open problems remain: How can

^{*}Electronic mail: bluman@math.ubc.ca

[†]Electronic mail: yangzz@math.ubc.ca

one further systematically extend a tree of nonlocally related PDE systems for a given PDE system and, of particular importance, if the given system has no local conservation laws and no known nonlocally related subsystem?

In this paper, we present a new systematic method for constructing nonlocally related PDE systems for a given PDE system via point symmetries. In particular, we will show that a nonlocally related system arises naturally from each point symmetry of a PDE system. It will be seen that this method can be considered as a generalization of the method for constructing nonlocally related subsystems. As a consequence, one is able to further extend the known procedure for the construction of trees of nonlocally related PDE systems. For a given PDE system, we also show that nonlocally related PDE systems arising from its point symmetries can also yield nonlocal symmetries. We show that if one applies invertible point transformations to the variables of a given PDE system, then one is able to obtain more nonlocally related subsystems. It will be seen that such an invertible point transformation does yield a nonlocally related subsystem if it is also a point symmetry of the given system.

This paper is organized as follows. In Sect. 2, we introduce the new systematic method to construct nonlocally related PDE systems. In Sect. 3, the new method is applied to the existence problem for subsystems. In Sect. 4, the new method is used to construct nonlocally related systems for nonlinear diffusion equations, nonlinear wave equations and nonlinear reaction-diffusion equations. In Sect. 5, the calculation of point symmetries for the nonlocally related systems constructed in Sect. 4, is shown to yield nonlocal symmetries for the considered example equations. Finally, in Sect. 6, the new results in this paper are summarized and open problems are posed.

In this work, we use the package GeM for Maple [2] for symmetry and conservation law analysis.

2. New method: nonlocally related PDE systems arising from point symmetries

Notation: Throughout this paper, in the situation of n independent variables $x = (x^1, \dots, x^n)$ and m dependent variables $u = (u^1, \dots, u^m)$, Let

$$\partial^p u = \left\{ \frac{\partial^p u^\mu}{\partial x^{i_1} \dots \partial x^{i_p}} \mid i_1, \dots, i_p = 1, \dots, n; \mu = 1, \dots, m \right\}$$

denote the set of all partial derivatives of order p and $\partial u \equiv \partial^1 u$.

Consider a scalar PDE $\mathbf{R}\{x, t; u\}$ with one dependent variable u and two independent variables (x, t) :

$$F(x, t, u, u_t, u_x, \partial^2 u, \dots, \partial^k u) = 0. \quad (2.1)$$

A systematic method to construct nonlocally related PDE systems of the scalar PDE $\mathbf{R}\{x, t; u\}$ (2.1) is presented in [3]. Here, the starting point is to use a local conservation law of the scalar PDE $\mathbf{R}\{x, t; u\}$ (2.1):

$$D_t \Phi(x, t, u, u_t, u_x, \partial^2 u, \dots, \partial^l u) + D_x \Psi(x, t, u, u_t, u_x, \partial^2 u, \dots, \partial^l u) = 0. \quad (2.2)$$

Based on the conservation law (2.2), one constructs a corresponding nonlocally related PDE system (potential system) $\mathbf{S}\{x, t; u, v\}$ given by

$$\begin{aligned} v_x &= \Phi(x, t, u, u_t, u_x, \partial^2 u, \dots, \partial^l u), \\ v_t &= -\Psi(x, t, u, u_t, u_x, \partial^2 u, \dots, \partial^l u), \\ F(x, t, u, u_t, u_x, \partial^2 u, \dots, \partial^k u) &= 0. \end{aligned} \quad (2.3)$$

In this paper, instead of using a conservation law as the starting point, we present a new systematic method to construct nonlocally related PDE system through the use of a point symmetry as the starting point.

Suppose a scalar PDE $\mathbf{R}\{x, t; u\}$ (2.1) has a point symmetry with infinitesimal generator $\mathbf{X}$. By introducing canonical coordinates $X = X(x, t, u)$, $T = T(x, t, u)$ and $U = U(x, t, u)$, one is able to map the infinitesimal generator $\mathbf{X}$ into the canonical form $\mathbf{Y} = \frac{\partial}{\partial U}$ corresponding to the invariance of PDE $\mathbf{R}\{x, t; u\}$ (2.1) under translation in U . Consequently, the scalar PDE $\mathbf{R}\{x, t; u\}$ (2.1) becomes the invertibly equivalent scalar PDE $\hat{\mathbf{R}}\{X, T; U\}$ in terms of the canonical coordinates (X, T, U) . The PDE $\hat{\mathbf{R}}\{X, T; U\}$ is of the form

$$\hat{F}(X, T, U_T, U_X, \dots, \partial^k U) = 0. \quad (2.4)$$

Introducing new variables α and β , related to the first partial derivatives of U , one obtains the PDE system $\tilde{\mathbf{R}}\{X, T; U, \alpha, \beta\}$ given by

$$\begin{aligned} \alpha &= U_T, \\ \beta &= U_X, \\ \tilde{F}(X, T, \alpha, \beta, \dots, \partial^{k-1} \alpha, \partial^{k-1} \beta) &= 0, \end{aligned} \quad (2.5)$$

where $\tilde{F}(X, T, \alpha, \beta, \dots, \partial^{k-1} \alpha, \partial^{k-1} \beta)$ is obtained from $\hat{F}(X, T, U_T, U_X, \dots, \partial^k U)$ after making the appropriate substitutions.

By construction, the PDE system $\tilde{\mathbf{R}}\{X, T; U, \alpha, \beta\}$ (2.5) is equivalent and locally related to the scalar PDE $\hat{\mathbf{R}}\{X, T; U\}$ (2.4), and hence equivalent and locally related to the given scalar PDE $\mathbf{R}\{x, t; u\}$ (2.1).

Excluding the dependent variable U from the PDE system $\tilde{\mathbf{R}}\{X, T; U, \alpha, \beta\}$ (2.5), one obtains the PDE system $\check{\mathbf{R}}\{X, T; \alpha, \beta\}$

$$\begin{aligned} \alpha_X &= \beta_T, \\ \check{F}(X, T, \alpha, \beta, \dots, \partial^{k-1} \alpha, \partial^{k-1} \beta) &= 0. \end{aligned} \quad (2.6)$$

The PDE system $\check{\mathbf{R}}\{X, T; \alpha, \beta\}$ (2.6) is nonlocally related and equivalent to the PDE $\hat{\mathbf{R}}\{X, T; U\}$. In particular, if $U = \Theta(X, T)$ solves the PDE $\hat{\mathbf{R}}\{X, T; U\}$ (2.4), then $(\alpha, \beta) = \left(\frac{\partial \Theta(X, T)}{\partial T}, \frac{\partial \Theta(X, T)}{\partial X}\right)$ is a solution of the PDE system $\check{\mathbf{R}}\{X, T; \alpha, \beta\}$ (2.6). Conversely, if $(\alpha, \beta) = (\Xi(X, T), \Omega(X, T))$ solves the PDE system $\check{\mathbf{R}}\{X, T; \alpha, \beta\}$ (2.6), due to the integrability relation $U_{TX} = U_{XT}$, there is a corresponding solution $U = \Theta(X, T)$ of the PDE $\hat{\mathbf{R}}\{X, T; U\}$. Moreover, the corresponding solution of the PDE $\hat{\mathbf{R}}\{X, T; U\}$ (2.4) is unique to within an arbitrary constant, i.e., if $U = \Theta(X, T)$ solves the PDE $\hat{\mathbf{R}}\{X, T; U\}$ (2.4), so does $U = \Theta(X, T) + C$ for any constant C . Moreover, since the scalar PDE $\hat{\mathbf{R}}\{X, T; U\}$ (2.4) and the given scalar PDE $\mathbf{R}\{x, t; u\}$ (2.1) are invertibly related, one concludes that the PDE system $\check{\mathbf{R}}\{X, T; \alpha, \beta\}$ (2.6) is a nonlocally related PDE system of the given PDE $\mathbf{R}\{x, t; u\}$ (2.1). Hence, the following theorem has been proved.

Theorem 2.1. Any point symmetry of a given scalar PDE $\mathbf{R}\{x, t; u\}$ (2.1) yields a nonlocally related equivalent PDE system of $\mathbf{R}\{x, t; u\}$ (2.1) given by the PDE system (2.6).

Corollary 2.1. Consider a scalar PDE $\mathbf{R}\{x, t; u\}$ given by

$$u_t = F(x, t, u_1, \dots, u_n), \quad (2.7)$$

where $u_n \equiv \frac{\partial^n u}{\partial x^n}$. Let $\beta = u_x$. Then the scalar PDE

$$\beta_t = D_x F(x, t, \beta, \dots, \beta_{n-1}) \quad (2.8)$$

is nonlocally related to the PDE $\mathbf{R}\{x, t; u\}$ (2.7).

Proof. Introducing the new variables α and β , related to the first partial derivatives of u , one obtains the following locally related PDE system $\tilde{\mathbf{R}}\{x, t; u, \alpha, \beta\}$ of the given PDE $\mathbf{R}\{x, t; u\}$ (2.7):

$$\begin{aligned} \alpha &= u_t, \\ \beta &= u_x, \\ \alpha &= F(x, t, \beta, \dots, \beta_{n-1}). \end{aligned} \quad (2.9)$$

Excluding the dependent variable u from the PDE system $\tilde{\mathbf{R}}\{x, t; u, \alpha, \beta\}$ (2.9), one obtains the PDE system $\check{\mathbf{R}}\{x, t; \alpha, \beta\}$

$$\begin{aligned} \alpha_x &= \beta_t, \\ \alpha &= F(x, t, \beta, \dots, \beta_{n-1}). \end{aligned} \quad (2.10)$$

From the previous discussion, the PDE system $\check{\mathbf{R}}\{x, t; \alpha, \beta\}$ (2.10) is nonlocally related to (2.7). Furthermore, one can exclude the dependent variable α from the PDE system $\check{\mathbf{R}}\{x, t; \alpha, \beta\}$ (2.10) to obtain the scalar PDE $\mathbf{R}\{x, t; \beta\}$

$$\beta_t = D_x F(x, t, \beta, \dots, \beta_{n-1}). \quad (2.11)$$

Since the excluded variable α can be expressed from the equations of the PDE system $\check{\mathbf{R}}\{x, t; \alpha, \beta\}$ (2.10) in terms of β and its derivatives, the PDE $\mathbf{R}\{x, t; \beta\}$ (2.11) is locally related to the PDE system $\check{\mathbf{R}}\{x, t; \alpha, \beta\}$ (2.10). Hence, the PDE $\mathbf{R}\{x, t; \beta\}$ (2.11) is nonlocally related and equivalent to the given PDE $\mathbf{R}\{x, t; u\}$ (2.7). $\square$

Note 1.1. *The situation for a PDE system with at least two dependent variables:* As presented above, the effective starting point of this method is that the given PDE is invariant under translation of its dependent variable. Now suppose a PDE system with at least two dependent variables has a point symmetry. Introducing canonical coordinates, one can always rewrite the given PDE system in a form that is invariant under translation of one of its dependent variables. Thus, in principle, one can apply this method to any PDE system that admits a point symmetry. Details will be presented in the next section.

Note 1.2. *Connection between our new method and the known potential systems method.* Our new method does not require the existence of a local conservation law of a given PDE system. Thus the new method is complementary to the previously known method for constructing nonlocally related PDE systems, especially in the case when a given PDE system is a scalar PDE.

Note 1.3. *The situation for a PDE system with at least three independent variables:* Our method can be adapted to a PDE system which has at least three independent variables. Without loss of generality, consider a scalar PDE $\mathbf{R}\{x; u\}$ with one dependent variable u and n independent variables $x = (x^1, \dots, x^n)$:

$$F(x, u, \partial u, \partial^2 u, \dots, \partial^k u) = 0. \quad (2.12)$$

Suppose the scalar PDE $\mathbf{R}\{x; u\}$ (2.12) has a point symmetry with infinitesimal generator $\mathbf{X}$. Introducing canonical coordinates $X^i = X^i(x^1, \dots, x^n, u)$, $i = 1, \dots, n$, and $U = U(x^1, \dots, x^n, u)$, one can map the scalar PDE $\mathbf{R}\{x; u\}$ (2.12) to an invertibly equivalent and hence locally related PDE $\hat{\mathbf{R}}\{X; U\}$ ($X = (X^1, \dots, X^n)$) of the form

$$\hat{F}(X, U_{X^1}, \dots, U_{X^n}, \partial^2 U, \dots, \partial^k U) = 0. \quad (2.13)$$

Introducing the new variables $\alpha = (\alpha^1, \dots, \alpha^n)$, related to the first partial derivatives of U , one obtains the PDE system $\tilde{\mathbf{R}}\{X; U, \alpha\}$ given by

$$\begin{aligned} \alpha^i &= U_{X^i}, \quad i = 1, \dots, n, \\ \tilde{F}(X, \alpha, \partial \alpha, \dots, \partial^{k-1} \alpha) &= 0, \end{aligned} \quad (2.14)$$

where $\tilde{F}(X, \alpha, \partial \alpha, \dots, \partial^{k-1} \alpha)$ is obtained from $\hat{F}(X, U_{X^1}, \dots, U_{X^n}, \partial^2 U, \dots, \partial^k U)$ after making the appropriate substitutions. The PDE system (2.14) is locally related to the PDE (2.13). Excluding U from the PDE system (2.14), one obtains the PDE system

$$\begin{aligned} \alpha_{X^j}^i - \alpha_{X^i}^j &= 0, \quad i, j = 1, \dots, n, \\ \tilde{F}(X, \alpha, \partial \alpha, \dots, \partial^{k-1} \alpha) &= 0. \end{aligned} \quad (2.15)$$

By construction, the PDE system (2.15) is nonlocally related to the scalar PDE (2.13), hence nonlocally related to the scalar PDE (2.12). Moreover, since the PDE system (2.15) has curl-type conservation laws, it is possible to yield nonlocal symmetries of the scalar PDE (2.12) from the PDE system (2.15) [1, 4, 5].

3. Subsystems arising from invertibly related PDE systems of a given PDE system

In [1] and references therein, it is shown that an important way of obtaining a nonlocally related system for a given PDE system with at least two dependent variables is through consideration of subsystems of the given PDE system. In particular, subsystems can arise directly through the elimination of one or more of the dependent variables of the given PDE system as well as indirectly through the elimination of one or more of the resulting dependent variables following a point transformation that involves an interchange of one or more of the dependent and independent variables of the given PDE system.

More generally, in principle any invertible point transformation of the dependent and independent variables of a given PDE system can be used to exclude one or more resulting dependent variables to obtain additional nonlocally related subsystems. However, often such an invertible point transformation does not lead to the exclusion of one or more dependent variables so that no subsystem is obtained. Moreover it can often happen that a resulting subsystem is not locally related to the given system.

Most importantly, the new method outlined in Sect. 2, indicates that if an invertible point transformation of the dependent and independent variables of a given PDE system is also a point symmetry of the given PDE system, then one always obtains, constructively, a nonlocally related system of the given PDE system. This is summarized in the following theorem.

Theorem 3.1. Consider a PDE system $\mathbf{R}\{x, t; u\}$ with two independent variables (x, t) and l dependent variables $u = (u^1, \dots, u^l)$, $l \geq 2$. Suppose $\mathbf{R}\{x, t; u\}$ has a point symmetry. In terms of its canonical coordinates, the point symmetry of $\mathbf{R}\{x, t; u\}$ is an invertible point transformation that maps $\mathbf{R}\{x, t; u\}$ to a locally related PDE system $\mathbf{S}\{X, T; U\}$ that has an explicit dependence on at most $l - 1$ dependent variables. A nonlocally related system of the PDE system $\mathbf{R}\{x, t; u\}$ arises naturally from the PDE system $\mathbf{S}\{X, T; U\}$.

Proof. Suppose the PDE system $\mathbf{R}\{x, t; u\}$ has a point symmetry with infinitesimal generator

$$\mathbf{X} = \xi(x, t, u) \frac{\partial}{\partial x} + \tau(x, t, u) \frac{\partial}{\partial t} + \eta^\nu(x, t, u) \frac{\partial}{\partial u^\nu}. \quad (3.1)$$

Let $\mathcal{F}$ be an invertible point transformation acting on (x, t, u) -space, which is given by

$$\begin{aligned} X &= f(x, t, u), \\ T &= g(x, t, u), \\ U^\nu &= \psi^\nu(x, t, u), \quad \nu = 1, \dots, l. \end{aligned} \quad (3.2)$$

Then $\mathcal{F}$ sends the PDE system $\mathbf{R}\{x, t; u\}$ to the PDE system $\mathbf{S}\{X, T; U\}$ and the infinitesimal generator (3.1) to the infinitesimal generator $\mathcal{F}_*(\mathbf{X})$, where $\mathcal{F}_*(\mathbf{X})$ is given by

$$\mathcal{F}_*(\mathbf{X}) = (\xi f_x + \tau f_t + \eta^\nu f_{u^\nu}) \frac{\partial}{\partial X} + (\xi g_x + \tau g_t + \eta^\nu g_{u^\nu}) \frac{\partial}{\partial T} + (\xi \psi_x^\rho + \tau \psi_t^\rho + \eta^\nu \psi_{u^\nu}^\rho) \frac{\partial}{\partial U^\rho}. \quad (3.3)$$

Since $\mathcal{F}$ is an invertible point transformation, $\mathcal{F}_*(\mathbf{X})$ is a point symmetry of PDE system $\mathbf{S}\{X, T; U\}$. One can now choose the transformation (3.2) to correspond to canonical coordinates for the point symmetry (3.1). This is accomplished by setting

$$\begin{aligned} \xi f_x + \tau f_t + \eta^\nu f_{u^\nu} &= 0, \\ \xi g_x + \tau g_t + \eta^\nu g_{u^\nu} &= 0, \\ \xi \psi_x^\rho + \tau \psi_t^\rho + \eta^\nu \psi_{u^\nu}^\rho &= 0, \quad \rho = 1, \dots, l - 1, \\ \xi \psi_x^l + \tau \psi_t^l + \eta^\nu \psi_{u^\nu}^l &= 1 \end{aligned} \quad (3.4)$$

so that $\mathcal{F}_*(\mathbf{X}) = \frac{\partial}{\partial U^l}$, which implies that PDE system $\mathbf{S}\{X, T; U\}$ is invariant under translation in U^l . Consequently, one can apply the procedure introduced in Sect. 2 to PDE system $\mathbf{S}\{X, T; U\}$ to generate a subsystem. Since the linear PDE system (3.4) always has nontrivial solutions, one is able to find a family of invertible point transformations that maps $\mathbf{R}\{x, t; u\}$ to a PDE system $\mathbf{S}\{X, T; U\}$ which yields a nonlocally related system from the exclusion in PDE system $\mathbf{S}\{X, T; U\}$ of an explicit dependence on the translated canonical variable U^l that arises from the solution of system (3.4).

Without loss of generality, we complete the proof for the case when $l = 2$. In this case, the PDE system $\mathbf{S}\{X, T; U\}$ can be rewritten as

$$F^\sigma(X, T, U^1, U_T^1, U_X^1, U_T^2, U_X^2, \dots) = 0, \quad \sigma = 1, 2, \dots, s. \quad (3.5)$$

Letting $\alpha = U_T^2$ and $\beta = U_X^2$, one obtains the locally related PDE system $\hat{\mathbf{S}}\{X, T; U^1, U^2, \alpha, \beta\}$ of the PDE system (3.5) given by

$$\begin{aligned}\alpha &= U_T^2, \\ \beta &= U_X^2, \\ F^\sigma(X, T, U^1, U_T^1, U_X^1, \alpha, \beta, \dots) &= 0, \quad \sigma = 1, 2, \dots, s.\end{aligned}\tag{3.6}$$

Excluding the dependent variable U^2 from the PDE system (3.6), one obtains the PDE system $\tilde{\mathbf{S}}\{X, T; U^1, \alpha, \beta\}$ given by

$$\begin{aligned}\alpha_X &= \beta_T, \\ F^\sigma(X, T, U^1, U_T^1, U_X^1, \alpha, \beta, \dots) &= 0, \quad \sigma = 1, 2, \dots, s,\end{aligned}\tag{3.7}$$

which is nonlocally related to the PDE system (3.5) $\square$

4. Examples of nonlocally related PDE systems arising from point symmetries

Example 4.1. Consider the scalar nonlinear diffusion equation

$$v_t = K(v_x) v_{xx},\tag{4.1}$$

where $K(v_x)$ is an arbitrary constitutive function.

Firstly, from its invariance under translations of its dependent variable v , one can apply directly the method presented in Sect. 2 to equation (4.1). Letting $u = v_x$, $f = v_t$, one obtains the corresponding locally related PDE system of (4.1) given by

$$\begin{aligned}u &= v_x, \\ f &= v_t, \\ f &= K(u)u_x.\end{aligned}\tag{4.2}$$

Excluding v from the PDE system (4.2), one obtains the nonlocally related PDE system of (4.1) given by

$$\begin{aligned}u_t &= f_x, \\ f &= K(u)u_x.\end{aligned}\tag{4.3}$$

Excluding f from the PDE system (4.3), one finds that a nonlocally related PDE of (4.1) is given by the nonlinear heat equation

$$u_t = (K(u)u_x)_x.\tag{4.4}$$

Secondly, since equation (4.1) is invariant under translations of its independent variable x , one can interchange x and v to get an invertibly related PDE of the PDE (4.1) given by

$$x_t = \frac{K\left(\frac{1}{x_v}\right) x_{vv}}{x_v^2}.\tag{4.5}$$

Introducing new variables $w = x_v$ and $g = x_t$, one obtains the locally related PDE system of the equation (4.1) given by

$$\begin{aligned}w &= x_v, \\ g &= x_t, \\ g &= \frac{K\left(\frac{1}{w}\right)w_v}{w^2}.\end{aligned}\tag{4.6}$$

Excluding x from the PDE system (4.6), one obtains the nonlocally related PDE system of the equation (4.1) given by

$$\begin{aligned} w_t &= g_v, \\ g &= \frac{K\left(\frac{1}{w}\right)w_v}{w^2}. \end{aligned} \quad (4.7)$$

Moreover, one can exclude the variable g from the PDE system (4.7) to get the locally related PDE of the PDE system (4.7) given by

$$w_t = \left(\frac{K\left(\frac{1}{w}\right)w_v}{w^2} \right)_v, \quad (4.8)$$

which in turn is nonlocally related to the nonlinear diffusion equation (4.1).

Thirdly, since equation (4.1) is invariant under translations of its independent variable t , one can interchange t and v to get an invertibly related PDE of the PDE (4.1) given by

$$t_v^2 - 2K\left(-\frac{t_x}{t_v}\right)t_x t_{xv} t_v + K\left(-\frac{t_x}{t_v}\right)t_x^2 t_{vv} + K\left(-\frac{t_x}{t_v}\right)t_{xx} t_v^2 = 0. \quad (4.9)$$

Introducing new variables $\alpha = t_v$ and $\beta = t_x$, one obtains the locally related PDE system of the equation (4.1) given by

$$\begin{aligned} \alpha &= t_v, \\ \beta &= t_x, \\ \alpha^2 - 2K\left(-\frac{\beta}{\alpha}\right)\beta\alpha_x\alpha + K\left(-\frac{\beta}{\alpha}\right)\beta^2\alpha_v + K\left(-\frac{\beta}{\alpha}\right)\beta_x\alpha^2 &= 0. \end{aligned} \quad (4.10)$$

Excluding t from the PDE system (4.10), one obtains the nonlocally related PDE system of the equation (4.1) given by

$$\begin{aligned} \alpha_x &= \beta_v, \\ \alpha^2 - 2K\left(-\frac{\beta}{\alpha}\right)\beta\alpha_x\alpha + K\left(-\frac{\beta}{\alpha}\right)\beta^2\alpha_v + K\left(-\frac{\beta}{\alpha}\right)\beta_x\alpha^2 &= 0. \end{aligned} \quad (4.11)$$

Finally, since equation (4.1) is invariant under the scaling symmetry generated by $\mathbf{X} = x\frac{\partial}{\partial x} + 2t\frac{\partial}{\partial t} + v\frac{\partial}{\partial v}$, one can use the corresponding canonical coordinates transformation given by

$$\begin{aligned} X &= \frac{t}{x^2}, \\ T &= \frac{v}{x}, \\ V &= \ln x \end{aligned} \quad (4.12)$$

to map the nonlinear diffusion equation (4.1) to the invertibly related PDE

$$\begin{aligned} -V_X V_T^2 - 4K\left(\frac{1+TV_T+2XV_X}{V_T}\right)XV_{TX}V_T + K\left(\frac{1+TV_T+2XV_X}{V_T}\right)V_{TT} + 4K\left(\frac{1+TV_T+2XV_X}{V_T}\right)V_{TT}XV_X \\ -K\left(\frac{1+TV_T+2XV_X}{V_T}\right)V_T^2 - 8K\left(\frac{1+TV_T+2XV_X}{V_T}\right)X^2V_XV_{TX}V_T + 4K\left(\frac{1+TV_T+2XV_X}{V_T}\right)X^2V_X^2V_{TT} \\ + 2K\left(\frac{1+TV_T+2XV_X}{V_T}\right)XV_XV_T^2 + 4K\left(\frac{1+TV_T+2XV_X}{V_T}\right)X^2V_T^2V_{XX} = 0. \end{aligned} \quad (4.13)$$

Introducing new variables $\varphi = V_T$ and $\psi = V_X$, one obtains the locally related PDE system of the equation (4.1) given by

$$\begin{aligned} \varphi &= V_T, \\ \psi &= V_X, \\ -\psi\varphi^2 - 4K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X\varphi_X\varphi + K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)\varphi_T + 4K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X\varphi_T\psi \\ -K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)\varphi^2 - 8K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X^2\psi\varphi_X\varphi + 4K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X^2\psi^2\varphi_T \\ + 2K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X\psi\varphi^2 + 4K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X^2\varphi^2\psi_X &= 0. \end{aligned} \quad (4.14)$$

Excluding V from the PDE system (4.15), one obtains the nonlocally related PDE system of the equation (4.1) given by

$$\begin{aligned} \varphi_X &= \psi_T, \\ -\psi\varphi^2 - 4K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X\varphi_X\varphi + K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)\varphi_T + 4K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X\varphi_T\psi \\ &- K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)\varphi^2 - 8K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X^2\psi\varphi_X\varphi + 4K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X^2\psi^2\varphi_T \\ &+ 2K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X\psi\varphi^2 + 4K\left(\frac{1+T\varphi+2X\psi}{\varphi}\right)X^2\varphi^2\psi_X = 0. \end{aligned} \quad (4.15)$$

Now take as the given PDE the nonlinear heat equation (4.4). From its invariance under translations in x , one can interchange x and u to obtain the locally related PDE

$$x_t = -\left(\frac{K(u)}{x_u}\right)_u. \quad (4.16)$$

Accordingly, let $p = x_u$ and $q = x_t$. Then one obtains the following locally related PDE of the nonlinear heat equation (4.4):

$$\begin{aligned} p &= x_u, \\ q &= x_t, \\ q &= -\left(\frac{K(u)}{p}\right)_u. \end{aligned} \quad (4.17)$$

Excluding the variable x from the PDE system (4.17), one obtains the nonlocally related PDE system of (4.17) given by

$$\begin{aligned} p_t &= q_u, \\ q &= -\left(\frac{K(u)}{p}\right)_u. \end{aligned} \quad (4.18)$$

Finally, after excluding the variable q from the PDE system (4.18), one obtains the locally related subsystem of (4.18) given by

$$p_t = -\left(\frac{K(u)}{p}\right)_{uu}. \quad (4.19)$$

By construction, the PDE (4.19) is nonlocally related to the nonlinear heat equation (4.4). The relationship between the constructed nonlocally related PDE systems is indicated in Figure 4.1.




Figure 4.1

Example 4.2. As a second example, consider the nonlinear wave equation

$$u_{tt} = (c^2(u)u_x)_x \quad (4.20)$$

with an arbitrary constitutive wave speed function $c(u)$.

In [6], a tree of nonlocally related systems was constructed for the nonlinear wave equation (4.20) through conservation law analysis and subsystems obtained by exclusions of dependent variables. We now show how to use point symmetries of the following potential system of the nonlinear wave equation (4.20)

$$\begin{aligned} v_x &= u_t, \\ v_t &= c^2(u)u_x \end{aligned} \quad (4.21)$$

to obtain further nonlocally related PDE systems of the nonlinear wave equation (4.20).

Due to its obvious invariance under translations in v and t , the PDE system (4.21) has a point symmetry with the infinitesimal generator $-\frac{\partial}{\partial t} + \frac{\partial}{\partial v}$. Corresponding canonical coordinates yield an invertible point transformation Ψ of the form :

$$\begin{aligned} X &= x, \\ T &= u, \\ U &= t + v, \\ V &= v. \end{aligned} \quad (4.22)$$

The transformation (4.22) maps the potential system (4.21) to the invertibly related PDE system

$$\begin{aligned} V_X U_T - V_T U_X - 1 &= 0, \\ V_T + c^2(T)U_X - c^2(T)V_X &= 0, \end{aligned} \quad (4.23)$$

which is invariant under translations in U and V .

First of all, by excluding the dependent variable V from the PDE system (4.23), one obtains its nonlocally related subsystem given by

$$\begin{aligned} c^4(T)U_{XX} - c^2(T)U_{XX}U_T^2 - c^2(T)U_{TT}U_X^2 - 2c^2(T)U_{TX} + 2c^2(T)U_{TX}U_TU_X \\ - 2c(T)c'(T)U_X + 2c(T)c'(T)U_X^2U_T + U_{TT} = 0, \end{aligned} \quad (4.24)$$

which, in turn, is a nonlocally related scalar PDE for the nonlinear wave equation (4.20).

Secondly, by excluding the dependent variable U from the PDE system (4.23), one obtains another nonlocally related subsystem given by

$$-2c(T)V_XV_TV_{TX} + c(T)V_{XX}V_T^2 - c^3(T)V_{XX} + c(T)V_X^2V_{TT} - 2c'(T)V_X^2V_T = 0, \quad (4.25)$$

which, in turn, is another nonlocally related scalar PDE for the nonlinear wave equation (4.13).

Note 4.1. After using the point (hodograph) transformation $x = x(u, v)$, $t = t(u, v)$, the given potential system (4.21) becomes the invertibly equivalent and thus locally related linear PDE system

$$\begin{aligned} t_u &= x_v, \\ t_v &= c^{-2}(u)x_u. \end{aligned} \quad (4.26)$$

Consequently, the linear wave equation

$$x_{vv} = \left(c^{-2}(u)x_u\right)_u \quad (4.27)$$

is obtained by excluding the dependent variable t from the PDE system (4.26). Now note that in the scalar PDE (4.25) $X = x$, $T = u$, $V = v$. Thus the invertible point transformation

$v = V, u = T, x = X$ maps (4.25) to the linear wave equation (4.27).

Example 4.3. As a third example, consider the nonlinear reaction-diffusion equation

$$u_t - u_{xx} = Q(u), \quad (4.28)$$

where the reaction term $Q(u)$ is an arbitrary constitutive function. For any nonlinear function $Q(u)$, the nonlinear reaction-diffusion equation (4.28) has no nontrivial local conservation law. Consequently, one cannot construct a nonlocally related PDE system of the nonlinear reaction-diffusion equation (4.28) via the conservation law-based method. In contrast, one can construct nonlocally related PDE systems of the scalar PDE (4.28) via the symmetry-based method introduced in this paper. The point symmetry classification of the nonlinear reaction-diffusion (4.28) is presented in [7] and exhibited in Table 4.1.

Table 4.1 Point symmetry classification for the nonlinear reaction-diffusion equation (4.28)

$Q(u)$	#	admitted point symmetries
arbitrary	2	$\mathbf{X}_1 = \frac{\partial}{\partial x}, \mathbf{X}_2 = \frac{\partial}{\partial t}$.
$u^a (a \neq 0, 1)$	3	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3 = u \frac{\partial}{\partial u} - (a-1)t \frac{\partial}{\partial t} - \frac{a-1}{2} x \frac{\partial}{\partial x}$.
e^u	3	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_4 = \frac{\partial}{\partial u} - t \frac{\partial}{\partial t} - \frac{1}{2} x \frac{\partial}{\partial x}$.
$u \ln u + bu$	4	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_5 = ue^t \frac{\partial}{\partial u}, \mathbf{X}_6 = 2e^t \frac{\partial}{\partial x} - xue^t \frac{\partial}{\partial u}$.

For arbitrary $Q(u)$, the nonlinear reaction-diffusion equation (4.28) has the exhibited two point symmetries: $\mathbf{X}_1$ and $\mathbf{X}_2$. Therefore, using the method introduced in Sect. 2 one can use interchanges of x and u and also t and u to construct two nonlocally related PDE systems of the nonlinear reaction-diffusion equation (4.28).

Firstly, after an interchange of the variables x and u , the nonlinear reaction-diffusion equation (4.28) becomes the invertibly related PDE given by

$$x_t = \frac{x_{uu} - Q(u)x_u^3}{x_u^2}. \quad (4.29)$$

Corresponding to the invariance of PDE (4.29) under translations of its dependent variable x , one obtains the following locally related PDE system for the nonlinear reaction-diffusion equation (4.28):

$$\begin{aligned} v &= x_u, \\ w &= x_t, \\ w &= \frac{v_u - Q(u)v^3}{v^2}. \end{aligned} \quad (4.30)$$

Excluding x from (4.30), one obtains the nonlocally related PDE system of the nonlinear reaction-diffusion equation (4.28) given by

$$\begin{aligned} v_t &= w_u, \\ w &= \frac{v_u - Q(u)v^3}{v^2}. \end{aligned} \quad (4.31)$$

In addition, one can exclude w from the PDE system (4.31) to get the scalar PDE

$$v_t = \left(\frac{v_u - Q(u)v^3}{v^2} \right)_u. \quad (4.32)$$

By construction, the scalar PDE (4.29) is locally related to the PDE system (4.31), and hence, nonlocally related to the nonlinear reaction-diffusion equation (4.28).

Secondly, after an interchange of the variables t and u , the nonlinear reaction-diffusion equation (4.28) becomes

$$t_u^2 + t_{xx}t_u^2 - 2t_xt_{ux}t_u + t_x^2t_{uu} - Q(u)t_u^3 = 0, \quad (4.33)$$

which is not in solved form and has mixed derivatives. Corresponding to the invariance of PDE (4.33) under translations of its dependent variable t , one introduces new variables $\alpha = t_x$ and $\beta = t_u$ to obtain the locally related PDE system of the nonlinear reaction-diffusion equation (4.28) given by

$$\begin{aligned} \alpha &= t_x, \\ \beta &= t_u, \\ \beta^2 + \alpha_x\beta^2 - 2\alpha\alpha_u\beta + \alpha^2\beta_u - Q(u)\beta^3 &= 0. \end{aligned} \quad (4.34)$$

Excluding t from PDE system (4.34), one obtains another nonlocally related PDE system of the nonlinear reaction-diffusion equation (4.28) given by

$$\begin{aligned} \alpha_u - \beta_x &= 0, \\ \beta^2 + \alpha_x\beta^2 - 2\alpha\alpha_u\beta + \alpha^2\beta_u - Q(u)\beta^3 &= 0. \end{aligned} \quad (4.35)$$

When $Q(u) = u^a$, ($a \neq 0, 1$), the nonlinear reaction-diffusion equation (4.28) admits one additional point symmetry $\mathbf{X}_3$. Without loss of generality, we consider the case $Q(u) = u^3$. Canonical coordinates induced by $\mathbf{X}_3$ are given by

$$\begin{aligned} X &= xu, \\ T &= \frac{t}{x^2}, \\ U &= -\ln x. \end{aligned} \quad (4.36)$$

In the (X, T, U) coordinates system, the corresponding nonlinear reaction-diffusion equation (4.28) becomes the invertibly related PDE

$$\begin{aligned} -U_T U_X^2 + 4T U_{TX} U_X + U_{XX} - 4U_{XX} T U_T - 3U_X^2 - 2X U_X^3 + 10T U_T U_X^2 \\ - 8T^2 U_T U_{TX} U_X + 4T^2 U_T^2 U_{XX} + 4T^2 U_X^2 U_{TT} - X^3 U_X^3 = 0. \end{aligned} \quad (4.37)$$

Accordingly, introducing the new variables $p = U_X$ and $q = U_T$, one obtains the locally related PDE system of the nonlinear reaction-diffusion equation (4.28) given by

$$\begin{aligned} p &= U_X, \\ q &= U_T, \\ -qp^2 + 4T q_X p + p_X - 4T p_X q - 3p^2 - 2X p^3 + 10T qp^2 - 8T^2 qq_X p \\ + 4T^2 q^2 p_X + 4T^2 p^2 q_T - X^3 p^3 &= 0. \end{aligned} \quad (4.38)$$

Excluding U from the PDE system (4.38), one obtains an additional nonlocally related PDE system of the corresponding nonlinear reaction-diffusion equation (4.28) given by

$$\begin{aligned} p_T &= q_X, \\ -qp^2 + 4T q_X p + p_X - 4T p_X q - 3p^2 - 2X p^3 + 10T qp^2 - 8T^2 qq_X p \\ + 4T^2 q^2 p_X + 4T^2 p^2 q_T - X^3 p^3 &= 0. \end{aligned} \quad (4.39)$$

When $Q(u) = e^u$, the nonlinear reaction-diffusion equation (4.28) admits one additional point symmetry $\mathbf{X}_4$. Canonical coordinates induced by $\mathbf{X}_4$ are given by

$$\begin{aligned} X &= u + 2 \ln x, \\ T &= \frac{t}{x^2}, \\ U &= -2 \ln x. \end{aligned} \quad (4.40)$$

In the (X, T, U) coordinates system, the corresponding nonlinear reaction-diffusion equation (4.28) becomes the invertibly related PDE

$$\begin{aligned} & -U_T U_X^2 - 2U_X^2 - 2U_X^3 + 6TU_X^2 U_T + 8TU_{TX} U_X + 4U_{XX} - 8TU_{XX} U_T \\ & - 8T^2 U_T U_{TX} U_X + 4T^2 U_T^2 U_{XX} + 4T^2 U_X^2 U_{TT} - e^X U_X^3 = 0. \end{aligned} \quad (4.41)$$

It follows that the introduction of the new variables $r = U_X$ and $s = U_T$ yields the locally related PDE system of the nonlinear reaction-diffusion equation (4.28) given by

$$\begin{aligned} & r = U_X, \\ & s = U_T, \\ & -sr^2 - 2r^2 - 2r^3 + 6Tr^2 s + 8Ts_X r + 4r_X - 8Tr_X s - 8T^2 ss_X r \\ & + 4T^2 s^2 r_X + 4T^2 r^2 s_T - e^X r^3 = 0. \end{aligned} \quad (4.42)$$

Excluding U from the PDE system (4.42), one obtains a third nonlocally related PDE system of the corresponding nonlinear reaction-diffusion (4.28) given by

$$\begin{aligned} & r_T = s_X, \\ & -sr^2 - 2r^2 - 2r^3 + 6Tr^2 s + 8Ts_X r + 4r_X - 8Tr_X s - 8T^2 ss_X r \\ & + 4T^2 s^2 r_X + 4T^2 r^2 s_T - e^X r^3 = 0. \end{aligned} \quad (4.43)$$

When $Q(u) = u \ln u + bu$, the nonlinear reaction-diffusion equation (4.28) has two additional point symmetries $\mathbf{X}_5$ and $\mathbf{X}_6$. Without loss of generality, we consider the case of $b = 1$, i.e., $Q(u) = u \ln u + u$.

Canonical coordinates induced by $\mathbf{X}_5$ are given by

$$\begin{aligned} & X = x, \\ & T = t, \\ & U = e^{-t} \ln u. \end{aligned} \quad (4.44)$$

In the (X, T, U) coordinates system, the corresponding nonlinear reaction-diffusion equation (4.28) becomes

$$U_T = U_{XX} + e^T U_X^2 + e^{-T}. \quad (4.45)$$

Thus one introduces the new variables $\zeta = U_X$ and $\theta = U_T$ to obtain the locally related PDE system of the nonlinear reaction-diffusion equation (4.28) given by

$$\begin{aligned} & \zeta = U_X, \\ & \theta = U_T, \\ & \theta = \zeta_X + e^T \zeta^2 + e^{-T}. \end{aligned} \quad (4.46)$$

Accordingly, one can exclude U and θ from PDE system (4.42) to obtain the nonlocally related scalar PDE of the corresponding nonlinear reaction-diffusion (4.28) given by

$$\zeta_T = \zeta_{XX} + 2e^T \zeta \zeta_X. \quad (4.47)$$

Canonical coordinates induced by $\mathbf{X}_6$ are given by

$$\begin{aligned} & X = e^{\frac{x^2}{4}} u, \\ & T = t, \\ & U = \frac{1}{2} e^{-t} x. \end{aligned} \quad (4.48)$$

In the (X, T, U) coordinates system, the corresponding nonlinear reaction-diffusion equation (4.28) becomes

$$U_T = \frac{e^{-2T} U_{XX} - 2X U_X^3 - 4X \ln X U_X^3}{4U_X^2}. \quad (4.49)$$

Hence, introducing the variables $\varphi = U_X$ and $\psi = U_T$, one obtains the locally related PDE system of the corresponding nonlinear reaction-diffusion equation (4.28) given by

$$\begin{aligned} \varphi &= U_X, \\ \psi &= U_T, \\ \psi &= \frac{e^{-2T} \varphi_X - 2X \varphi^3 - 4X \ln X \varphi^3}{4\varphi^2}. \end{aligned} \quad (4.50)$$

Excluding U and ψ from the PDE system (4.42), one obtains another nonlocally related PDE system of the corresponding nonlinear reaction-diffusion (4.28) given by

$$\varphi_T = \left(\frac{e^{-2T} \varphi_X - 2X \varphi^3 - 4X \ln X \varphi^3}{4\varphi^2} \right)_X. \quad (4.51)$$

The relationship between the constructed nonlocally related PDE systems for the nonlinear reaction-diffusion equation is illustrated in Figure 4.2.




Figure 4.2

5. Nonlocal symmetries for examples presented in Section 4

A symmetry of a PDE system is any mapping of its solution manifold into itself, i.e., a symmetry maps any solution of a PDE system to a solution of the same system. Consequently, continuous symmetries of PDE systems are defined topologically and hence not restricted to just point or local symmetries. Thus, in principle, any nontrivial PDE system has symmetries. The problem is to find and use symmetries. Practically, to find a symmetry of a PDE system systematically, one is essentially restricted to transformations acting locally on some finite-dimensional space, whose variables include but are not restricted to just the independent and dependent variables of the PDE system. From this point of view, local symmetries, whose infinitesimals depend at most on a finite number of derivatives of the dependent variables of the given PDE system, constitute only a subset of the total set of symmetries of a PDE system. Otherwise, there exist nonlocal symmetries of a PDE system. When nonlocal symmetries can be found for a given PDE system, it may be possible to use such symmetries systematically to generate further exact solutions from its known solutions, to construct invariant solutions, to find linearizations, or find additional nonlocally related systems. However, when one directly applies Lie's algorithm to find nonlocal symmetries, the coefficients of the infinitesimal generators should essentially involve integrals of the dependent variables and their derivatives [8–10]. It is difficult to set up and obtain solutions of corresponding determining equations for such coefficients.

In [1] and references therein, systematic ways are presented to find nonlocal symmetries by applying Lie's algorithm to nonlocally related PDE systems. In the framework of such

nonlocally related PDE systems, nonlocal symmetries of a given PDE system $\mathbf{R}\{x, t; u\}$ can arise from point symmetries of any PDE system in a tree of nonlocally related systems that includes $\mathbf{R}\{x, t; u\}$. In particular, three different types of such symmetries can be found.

1. Nonlocal symmetries arising from point symmetries of potential systems of $\mathbf{R}\{x, t; u\}$.
2. Nonlocal symmetries arising from point symmetries of nonlocally related subsystems of $\mathbf{R}\{x, t; u\}$.
3. Nonlocal symmetries arising from point symmetries of nonlocally related subsystems of potential systems of $\mathbf{R}\{x, t; u\}$.

For Type 1, a point symmetry of a potential system of $\mathbf{R}\{x, t; u\}$ yields a nonlocal symmetry of $\mathbf{R}\{x, t; u\}$ if and only if the infinitesimal components corresponding to its given variables (x, t, u) involve nonlocal variables. For Types 2 and 3, one must trace back to see whether the obtained point symmetry yields a nonlocal symmetry of $\mathbf{R}\{x, t; u\}$.

In this section, we apply point symmetry analysis to the examples presented in Sect. 4 to seek nonlocal symmetries. A point symmetry classification analysis for all nonlocally related systems arising for the reaction-diffusion equation (4.2), i.e, Example 4.3, yields no nonlocal symmetries. In the following subsections, we list the corresponding nonlocal symmetries obtained for Examples 4.1 and 4.2.

5.1 Nonlocal symmetries for the nonlinear diffusion equation (4.1) and the nonlinear heat equation (4.4)

In [3], the point symmetry classification is given for the PDE system

$$\begin{aligned} v_x &= u, \\ v_t &= K(u)u_x, \end{aligned} \tag{5.1}$$

which is locally related to the nonlinear diffusion equation (4.1). Moreover, due to the invariance of PDE system (5.1) under translations in v , as shown in Sect. 2, the method presented in this paper systematically yields the nonlinear heat equation as a nonlocally related subsystem of (5.1) and hence as a nonlocally related system of the nonlinear diffusion equation (4.1). The point symmetry classification of system (5.1) is presented in Table 5.1, modulo the group of equivalence transformations of the class of PDEs (5.1).

Table 5.1 Point symmetry classification for the PDE system (5.1)

$K(u)$	#	admitted point symmetries
arbitrary	4	$\mathbf{X}_1 = \frac{\partial}{\partial x}, \mathbf{X}_2 = \frac{\partial}{\partial t}, \mathbf{X}_3 = x\frac{\partial}{\partial x} + 2t\frac{\partial}{\partial t} + v\frac{\partial}{\partial v}, \mathbf{X}_4 = \frac{\partial}{\partial v}.$
u^μ	5	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \mathbf{X}_4, \mathbf{X}_5 = x\frac{\partial}{\partial x} + \frac{2}{\mu}u\frac{\partial}{\partial u} + \left(1 + \frac{2}{\mu}\right)v\frac{\partial}{\partial v}.$
e^u	5	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \mathbf{X}_4, \mathbf{X}_6 = x\frac{\partial}{\partial x} + 2\frac{\partial}{\partial u} + (2x + v)\frac{\partial}{\partial v}.$
u^{-2}	∞	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \mathbf{X}_4, \mathbf{X}_5 (\mu = -2),$ $\mathbf{X}_7 = -xv\frac{\partial}{\partial x} + (xu + v)u\frac{\partial}{\partial u} + 2t\frac{\partial}{\partial v},$ $\mathbf{X}_8 = -x(2t + v^2)\frac{\partial}{\partial x} + 4t^2\frac{\partial}{\partial t} + u(6t + 2xuv + v^2)\frac{\partial}{\partial u} + 4tv\frac{\partial}{\partial v},$ $\mathbf{X}_\infty = A(v, t)\frac{\partial}{\partial x} - u^2B(v, t)\frac{\partial}{\partial u},$ where $(A(v, t), B(v, t))$ is an arbitrary solution of the linear system $A_t = B_v, A_v = B.$
$u^\mu(1 + bu)^{-(\mu+2)}$	5	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \mathbf{X}_4,$ $\mathbf{X}_9 = bv\frac{\partial}{\partial x} + \mu t\frac{\partial}{\partial t} - (1 + bu)u\frac{\partial}{\partial u} - v\frac{\partial}{\partial v}.$
$(1 + bu)^{-2}e^{u/(1+bu)}$	5	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \mathbf{X}_4,$ $\mathbf{X}_{10} = b(2x + bv)\frac{\partial}{\partial x} + (1 + 2b)t\frac{\partial}{\partial t} - (1 + bu)^2\frac{\partial}{\partial u} - x\frac{\partial}{\partial v}.$
$(u^2 + (1 + bu)^2)^{-1}e^{\lambda \arctan(u/(1+bu))}$	5	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \mathbf{X}_4,$ $\mathbf{X}_{11} = (2bx + (b^2 + 1)v)\frac{\partial}{\partial x} + (\lambda + 2b)t\frac{\partial}{\partial t} - ((1 + bu)^2 + u^2)\frac{\partial}{\partial u} - x\frac{\partial}{\partial v}.$
$(u^2 - (1 + bu)^2)^{-1}e^{\lambda \operatorname{arctanh}(u/(1+bu))}$	5	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \mathbf{X}_4,$ $\mathbf{X}_{12} = (2bx + (b^2 - 1)v)\frac{\partial}{\partial x} + (\lambda + 2b)t\frac{\partial}{\partial t} - ((1 + bu)^2 - u^2)\frac{\partial}{\partial u} - x\frac{\partial}{\partial v}.$

In [11], the point symmetry classification is given for the nonlinear heat equation (4.4). This classification, modulo the group of equivalence transformations of the class of PDEs (4.4), is presented in Table 5.2.

Table 5.2 Point symmetry classification for the nonlinear heat equation (4.4)

$K(u)$	#	admitted point symmetries
arbitrary	3	$\mathbf{Y}_1 = \frac{\partial}{\partial x}, \mathbf{Y}_2 = \frac{\partial}{\partial t}, \mathbf{Y}_3 = x\frac{\partial}{\partial x} + 2t\frac{\partial}{\partial t}.$
u^μ	4	$\mathbf{Y}_1, \mathbf{Y}_2, \mathbf{Y}_3, \mathbf{Y}_4 = x\frac{\partial}{\partial x} + \frac{2}{\mu}u\frac{\partial}{\partial u}.$
e^u	4	$\mathbf{Y}_1, \mathbf{Y}_2, \mathbf{Y}_3, \mathbf{Y}_5 = x\frac{\partial}{\partial x} + 2\frac{\partial}{\partial u}.$
$u^{-\frac{4}{3}}$	5	$\mathbf{Y}_1, \mathbf{Y}_2, \mathbf{Y}_3, \mathbf{Y}_4 (\mu = -\frac{4}{3}), \mathbf{Y}_6 = x^2\frac{\partial}{\partial x} - 3xu\frac{\partial}{\partial u}.$

Proposition 5.1. The symmetry $\mathbf{Y}_6$ yields a nonlocal symmetry of the nonlinear diffusion equation (4.1).

Proof. Suppose the symmetry $\mathbf{Y}_6$ yields a local symmetry of the nonlinear diffusion equation (4.1). Since the nonlinear diffusion equation (4.1) and the PDE system (5.1) are locally related, $\mathbf{Y}_6$ must also yield a local symmetry $\hat{\mathbf{Y}}_6$ of the PDE system (5.1). Consequently, there must exist a function $f([u], [v])$, where $[u]$ and $[v]$ denote that f depends on x, t, u, v and their derivatives, such that, in evolutionary form, $\hat{\mathbf{Y}}_6 = (-3xu - x^2u_x)\frac{\partial}{\partial u} + f([u], [v])\frac{\partial}{\partial v}$ is a local symmetry of the PDE system (5.1). Since $v_x = u, v_t = u^{-\frac{4}{3}}u_x$ and $u_t = (u^{-\frac{4}{3}}u_x)_x$, without loss of generality one can restrict $f([u], [v])$ to be of the form $f(x, t, u, v, u_x, u_{xx}, \dots)$ depending on x, t, u and the partial derivatives of u with respect to x . Firstly, suppose $f([u], [v])$ is of the

form $f(x, t, u, v, u_x)$. Applying $\hat{\mathbf{Y}}_6^{(\infty)}$ to the PDE system (5.1), one obtains

$$\begin{aligned} f_x + f_u u_x + f_v v_x + f_{u_x} u_{xx} &= -3xu - x^2 u_x, \\ f_t + f_u u_t + f_v v_t + f_{u_x} u_{tx} &= \frac{4}{3}(3xu + x^2 u_x)u^{-\frac{7}{3}}u_x + D_x(-3xu - x^2 u_x)u^{-\frac{4}{3}} \end{aligned} \quad (5.2)$$

on every solution of the PDE system (5.1). After making appropriate substitutions and equating the coefficients of the term u_{xx} , one obtains $f_{u_x} = 0$. By similar reasoning, one can show that $f(x, t, u, v, u_x, u_{xx}, \dots)$ has no dependence on any partial derivative of u with respect to x . Hence, $f([u], [v])$ is of the form $f(x, t, u, v)$. Consequently, if $\mathbf{Y}_6$ yields a local symmetry of the nonlinear diffusion equation (4.1), then $\hat{\mathbf{Y}}_6$ must be a point symmetry of the PDE system (5.1).

Comparing Tables 5.1 and 5.2, one immediately sees that symmetry $\mathbf{Y}_6$ does not yield a point symmetry of the PDE system (5.1). This follows from the fact that when $K(u) = u^{-\frac{4}{3}}$, the PDE system (5.1) has no point symmetry whose infinitesimal components corresponding to the variables (x, t) are the same as those for $\mathbf{Y}_6$. Hence, $\mathbf{Y}_6$ yields a nonlocal symmetry of the nonlinear diffusion equation (4.1). $\square$

Note 5.1. This example demonstrates that the nonlocally related PDE system obtained by the method presented in Sect. 2 can yield nonlocal symmetries of a given PDE system.

Now consider the nonlocally related class of scalar PDEs (4.8). The equivalence transformations for this class of PDEs arise from the six generators

$$\begin{aligned} \mathbf{E}_1 &= \frac{\partial}{\partial v}, & \mathbf{E}_2 &= \frac{\partial}{\partial w} + \frac{2K}{w} \frac{\partial}{\partial K}, & \mathbf{E}_3 &= w \frac{\partial}{\partial w} + 2K \frac{\partial}{\partial K}, \\ \mathbf{E}_4 &= v \frac{\partial}{\partial v} + 2K \frac{\partial}{\partial K}, & \mathbf{E}_5 &= t \frac{\partial}{\partial t} - K \frac{\partial}{\partial K}, & \mathbf{E}_6 &= \frac{\partial}{\partial t}. \end{aligned} \quad (5.3)$$

Thus the group of equivalence transformations for the class of PDEs (4.8) is given by

$$\begin{aligned} \tilde{v} &= a_3 v + a_1, \\ \tilde{t} &= a_5 t + a_6, \\ \tilde{w} &= a_4 w + a_2, \\ \tilde{K} &= \frac{a_3^2 a_4^2 (w + a_2)^2}{a_5 w^2} K, \end{aligned} \quad (5.4)$$

where $a_1, \dots, a_6$ are arbitrary constants with $a_3, a_4, a_5 \neq 0$.

In Table 5.3, we present the point symmetry classification of the PDE (4.8), modulo its group of equivalence transformations (5.4).

Table 5.3 Point symmetry classification for the PDE (4.8)

$K(1/w)$	$K(u)$	#	admitted point symmetries in (t, v, w) coordinates	admitted point symmetries in (t, v, u) coordinates
arbitrary	arbitrary	3	$\mathbf{Z}_1 = \frac{\partial}{\partial t}, \mathbf{Z}_2 = \frac{\partial}{\partial v}, \mathbf{Z}_3 = 2t \frac{\partial}{\partial t} + v \frac{\partial}{\partial v}.$	$\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z}_3.$
$w^{-\mu}$	u^μ	4	$\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z}_3,$ $\mathbf{Z}_4 = (2 + \mu)v \frac{\partial}{\partial v} - 2w \frac{\partial}{\partial w}.$	$\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z}_3.$ $\mathbf{Z}_4 = (2 + \mu)v \frac{\partial}{\partial v} + 2u \frac{\partial}{\partial u}.$
$w^{\frac{2}{3}}$	$u^{-\frac{2}{3}}$	5	$\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z}_3, \mathbf{Z}_4 (\mu = -\frac{2}{3}),$ $\mathbf{Z}_5 = 3vw \frac{\partial}{\partial w} - v^2 \frac{\partial}{\partial v}.$	$\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z}_3, \mathbf{Z}_4 (\mu = -\frac{2}{3}),$ $\mathbf{Z}_5 = -3uv \frac{\partial}{\partial u} - v^2 \frac{\partial}{\partial v}.$
w^2	u^{-2}	∞	$\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z}_3, \mathbf{Z}_4 (\mu = -2),$ $\mathbf{Z}_6 = vw \frac{\partial}{\partial w} - 2t \frac{\partial}{\partial v}.$ $\mathbf{Z}_7 = 4t^2 \frac{\partial}{\partial t} + 4vt \frac{\partial}{\partial v} - (2t + v^2)w \frac{\partial}{\partial w}.$ $\mathbf{Z}_\infty = F(t, v) \frac{\partial}{\partial w},$ where $F(t, v)$ satisfies $F_t = F_{vv}.$	$\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z}_3, \mathbf{Z}_4 (\mu = -2),$ $\mathbf{Z}_6 = -uv \frac{\partial}{\partial u} - 2t \frac{\partial}{\partial v}.$ $\mathbf{Z}_7 = 4t^2 \frac{\partial}{\partial t} + 4vt \frac{\partial}{\partial v} + (2t + v^2)u \frac{\partial}{\partial u}.$ $\mathbf{Z}_\infty = -u^2 F(t, v) \frac{\partial}{\partial u},$ where $F(t, v)$ satisfies $F_t = F_{vv}.$

By similar reasoning as in the proof of Proposition 5.1, one can show that if a point symmetry $\mathbf{Z}$ yields a local symmetry of the nonlinear diffusion equation (4.1), then $\mathbf{Z}$ must yield a point symmetry of the PDE system (5.1). Comparing the infinitesimal components for the variables (u, v) in Tables 5.1 and 5.3, one sees that when $K(u) = u^{-\frac{2}{3}}$, the point symmetry $\mathbf{Z}_5$ of the PDE (4.8) yields a nonlocal symmetry of the nonlinear diffusion equation (4.1).

Moreover, comparing Tables 5.2 and 5.3, one also sees that when $K(u) = u^{-\frac{2}{3}}$, since its infinitesimal component for the variable u has an essential dependence on the variable v , the symmetry $\mathbf{Z}_5$ of the PDE (4.8) yields a nonlocal symmetry of the nonlinear heat equation (4.4), which cannot be obtained through its potential system (5.1). By similar reasoning, when $K(u) = u^{-2}$, one can show that the symmetries $\mathbf{Z}_6$, $\mathbf{Z}_7$ and $\mathbf{Z}_\infty$ of the PDE (4.8) yield nonlocal symmetries of the nonlinear heat equation (4.4).

The equivalence transformations of the class of PDEs (4.19) arise from the six generators

$$\begin{aligned} \mathbf{E}_1 &= \frac{\partial}{\partial u}, & \mathbf{E}_2 &= u \frac{\partial}{\partial u} + 2K \frac{\partial}{\partial K}, & \mathbf{E}_3 &= p \frac{\partial}{\partial p} + 3K \frac{\partial}{\partial K}, \\ \mathbf{E}_4 &= t \frac{\partial}{\partial t} - K \frac{\partial}{\partial K}, & \mathbf{E}_5 &= \frac{\partial}{\partial t} & \mathbf{E}_6 &= u^2 \frac{\partial}{\partial u} - up \frac{\partial}{\partial p} - Ku \frac{\partial}{\partial K}. \end{aligned} \quad (5.5)$$

Hence, the five-parameter group of equivalence transformations of the PDE class (4.19), arising from the first five generators of (5.5), is given by

$$\begin{aligned} \tilde{u} &= a_2 u + a_1, \\ \tilde{t} &= a_4 t + a_5, \\ \tilde{p} &= a_3 p, \\ \tilde{K} &= \frac{a_2^2 a_3}{a_4} K, \end{aligned} \quad (5.6)$$

where $a_1, \dots, a_5$ are arbitrary constants with $a_2, a_3, a_4 \neq 0$.

The generator $\mathbf{E}_6$ yields the additional one-parameter group of equivalence transformations given by

$$\tilde{u} = \frac{u}{1 - a_6 u}, \quad \tilde{t} = t, \quad \tilde{p} = (1 - a_6 u)^3 p, \quad \tilde{K} = (1 - a_6 u)^5 K, \quad (5.7)$$

where a_6 is an arbitrary constant.

In Table 5.4, we present the point symmetry classification of the PDE (4.19), modulo its group of equivalence transformations.

Table 5.4 Point symmetry classification for the PDE (4.19)

$K(u)$	#	admitted point symmetries
arbitrary	2	$\mathbf{W}_1 = \frac{\partial}{\partial t}, \mathbf{W}_2 = 2t \frac{\partial}{\partial t} + p \frac{\partial}{\partial p}.$
u^μ	3	$\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_3 = 2u \frac{\partial}{\partial u} + (\mu - 2)p \frac{\partial}{\partial p}.$
e^u	3	$\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_4 = 2 \frac{\partial}{\partial u} + p \frac{\partial}{\partial p}.$
$u^\mu (1 + bu)^{-(\mu+2)}$	3	$\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_5 = 2u(1 + bu) \frac{\partial}{\partial u} - p(6bu - \mu + 2) \frac{\partial}{\partial p}.$
$(1 + bu)^{-2} e^{u/(1+bu)}$	3	$\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_6 = 2(1 + bu)^2 \frac{\partial}{\partial u} - p(6b^2 u + 6b - 1) \frac{\partial}{\partial p}.$
$\frac{1}{u^2 + (1+bu)^2} e^{\lambda \arctan(u/(1+bu))}$	3	$\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_7 = 2(u^2 + (1 + bu)^2) \frac{\partial}{\partial u} - p(6b^2 u + 6u + 6b - \lambda) \frac{\partial}{\partial p}.$
$\frac{1}{u^2 - (1+bu)^2} e^{\lambda \operatorname{arctanh}(u/(1+bu))}$	3	$\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_8 = 2((1 + bu)^2 - u^2) \frac{\partial}{\partial u} - p(6b^2 u - 6u + 6b - \lambda) \frac{\partial}{\partial p}.$
u^{-2}	4	$\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_3 (\mu = -2), \mathbf{W}_9 = u^2 \frac{\partial}{\partial u} - 3pu \frac{\partial}{\partial p}.$

Similar to the situation in Proposition 5.1, one can show that if a point symmetry $\mathbf{W}$ yields a local symmetry $\hat{\mathbf{W}}$ of the nonlinear heat equation (4.4), then $\hat{\mathbf{W}}$ must be a contact symmetry (which could be a point symmetry) of the nonlinear heat equation (4.4). In [12], it is shown that any contact symmetry of the nonlinear heat equation (4.4) is a point symmetry of (4.4). Comparing Tables 5.2 and 5.4, one immediately sees that the symmetry $\mathbf{W}_5$ yields a nonlocal symmetry of the nonlinear heat equation (4.4), since when $K(u) = u^\mu(1 + bu)^{-(\mu+2)}$, the nonlinear heat equation (4.4) has no point symmetry whose infinitesimal component corresponding to the variable u is the same as that for $\mathbf{W}_5$. By similar reasoning, the symmetries $\mathbf{W}_6$, $\mathbf{W}_7$, $\mathbf{W}_8$ and $\mathbf{W}_9$ also yield nonlocal symmetries of the nonlinear heat equation (4.4).

5.2 Nonlocal symmetries for the nonlinear wave equation (4.20)

In [13], the point symmetry classification is given for the class of nonlinear wave equations (4.20), which is presented in Table 5.5, modulo its group of equivalence transformations.

Table 5.5 Point symmetry classification for the nonlinear wave equation (4.20)

$c(u)$	#	admitted point symmetries
arbitrary	3	$\mathbf{X}_1 = \frac{\partial}{\partial x}, \mathbf{X}_2 = \frac{\partial}{\partial t}, \mathbf{X}_3 = x \frac{\partial}{\partial x} + t \frac{\partial}{\partial t}.$
u^μ	4	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \mathbf{X}_4 = \mu x \frac{\partial}{\partial x} + u \frac{\partial}{\partial u}.$
e^u	4	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \mathbf{X}_5 = x \frac{\partial}{\partial x} + \frac{\partial}{\partial u}.$
u^{-2}	5	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \mathbf{X}_4 (\mu = -2), \mathbf{X}_6 = t^2 \frac{\partial}{\partial t} + tu \frac{\partial}{\partial u}.$
$u^{-\frac{2}{3}}$	5	$\mathbf{X}_1, \mathbf{X}_2, \mathbf{X}_3, \mathbf{X}_4 (\mu = -\frac{2}{3}), \mathbf{X}_7 = x^2 \frac{\partial}{\partial x} - 3xu \frac{\partial}{\partial u}.$

In [14], the point symmetry classification is given for the potential system (4.21), which is exhibited in Table 5.6, modulo its group of equivalence transformations.

Table 5.6 Point symmetry classification for the PDE system (4.21)

$c(u)$	#	admitted point symmetries
arbitrary	∞	infinite number of point symmetries $\mathbf{Y}_\infty$ following from the invertibly related linear system after an interchange of independent and dependent variables.
arbitrary	4 more	$\mathbf{Y}_\infty, \mathbf{Y}_1 = \frac{\partial}{\partial t}, \mathbf{Y}_2 = \frac{\partial}{\partial x}, \mathbf{Y}_3 = \frac{\partial}{\partial v}, \mathbf{Y}_4 = x\frac{\partial}{\partial x} + t\frac{\partial}{\partial t}.$
$u^\mu (\mu \neq 0, -1)$	6 more	$\mathbf{Y}_\infty, \mathbf{Y}_1, \mathbf{Y}_2, \mathbf{Y}_3, \mathbf{Y}_4,$ $\mathbf{Y}_5 = \mu t \frac{\partial}{\partial t} - u \frac{\partial}{\partial u} - (1 + \mu)v \frac{\partial}{\partial v},$ $\mathbf{Y}_6 = -((2\mu + 1)tv + xu) \frac{\partial}{\partial t} - (tu^{1+2\mu} + xv) \frac{\partial}{\partial x} + 2uv \frac{\partial}{\partial u}$ $+ \left[(1 + \mu)v^2 + \frac{u^{2+2\mu}}{1+\mu} \right] \frac{\partial}{\partial v}.$
e^u	6 more	$\mathbf{Y}_\infty, \mathbf{Y}_1, \mathbf{Y}_2, \mathbf{Y}_3, \mathbf{Y}_4,$ $\mathbf{Y}_7 = x \frac{\partial}{\partial x} + \frac{\partial}{\partial u} + v \frac{\partial}{\partial v},$ $\mathbf{Y}_8 = -(2vt + x) \frac{\partial}{\partial t} - e^{2u}t \frac{\partial}{\partial x} + 2v \frac{\partial}{\partial u} + (e^{2u} + v^2) \frac{\partial}{\partial v}.$
u^{-1}	6 more	$\mathbf{Y}_\infty, \mathbf{Y}_1, \mathbf{Y}_2, \mathbf{Y}_3, \mathbf{Y}_4, \mathbf{Y}_5 (\mu = -1),$ $\mathbf{Y}_9 = (tv - xu) \frac{\partial}{\partial t} - (tu^{-1} + xv) \frac{\partial}{\partial x} + 2uv \frac{\partial}{\partial u} + 2 \ln u \frac{\partial}{\partial v}.$
$c(u)$ satisfies (a), (b) or (c): (a) $c' = c^2 \mu^{-1} \sinh(\mu \ln c)$ (b) $c' = c^2 \mu^{-1} \sin(\mu \ln c)$ (c) $c' = c^2 \mu^{-1} \cosh(\mu \ln c)$	6 more	$\mathbf{Y}_\infty, \mathbf{Y}_1, \mathbf{Y}_2, \mathbf{Y}_3, \mathbf{Y}_4,$ $\mathbf{Y}_{10,11} = e^{\pm v} \left\{ ((2 + \Gamma')t \pm \Gamma x) \frac{\partial}{\partial t} + (\Gamma'x \pm c^2 \Gamma t) \frac{\partial}{\partial x} - 2\Gamma \frac{\partial}{\partial u} \right.$ $\left. \mp 2(\Gamma' + 1) \frac{\partial}{\partial v} \right\},$ where $\Gamma = c/c'.$

The point symmetry classification of the PDE (4.27), modulo the equivalence transformations, is presented in [6].

The equivalence transformations for the PDE class (4.24) arise from the five generators

$$\begin{aligned} \mathbf{E}_1 &= \frac{\partial}{\partial T}, & \mathbf{E}_2 &= \frac{\partial}{\partial X}, & \mathbf{E}_3 &= \frac{\partial}{\partial U} \\ \mathbf{E}_4 &= T \frac{\partial}{\partial T} + X \frac{\partial}{\partial X} + U \frac{\partial}{\partial U}, & \mathbf{E}_5 &= -T \frac{\partial}{\partial T} + X \frac{\partial}{\partial X} + c \frac{\partial}{\partial c}. \end{aligned} \quad (5.8)$$

Correspondingly, the five-parameter group of equivalence transformations for the class of PDEs (4.19) is given by

$$\begin{aligned} \tilde{T} &= \frac{a_4}{a_5} T + a_1, \\ \tilde{X} &= a_4 a_5 X + a_2, \\ \tilde{U} &= a_4 U + a_3, \\ \tilde{c} &= a_5 c, \end{aligned} \quad (5.9)$$

where $a_1, \dots, a_5$ are arbitrary constants with $a_4, a_5 \neq 0$.

The point symmetry classification of the PDE (4.24), modulo its equivalence transformations, is presented in Table 5.7.

Table 5.7 Point symmetry classification for the PDE (4.24)

$c(T)$	$c(u)$	#	admitted point symmetries in (X, T, U) coordinates	admitted point symmetries in (x, t, u, v) coordinates
arbitrary	arbitrary	3	$\mathbf{W}_1 = \frac{\partial}{\partial U}, \mathbf{W}_2 = \frac{\partial}{\partial X},$ $\mathbf{W}_3 = (X + \int^T c^2(\xi)d\xi)\frac{\partial}{\partial X} + U\frac{\partial}{\partial U}.$	$\mathbf{W}_1 = \frac{\partial}{\partial t}, \mathbf{W}_2 = \frac{\partial}{\partial x},$ $\mathbf{W}_3 = (x + \int^u c^2(\xi)d\xi)\frac{\partial}{\partial x} + (t + v)\frac{\partial}{\partial t}.$
T^μ	u^μ	4	$\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_3 (c(T) = T^\mu),$ $\mathbf{W}_4 = T\frac{\partial}{\partial T} + (2\mu + 1)X\frac{\partial}{\partial X}$ $+ (\mu + 1)U\frac{\partial}{\partial U}.$	$\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_3 (c(u) = u^\mu),$ $\mathbf{W}_4 = u\frac{\partial}{\partial u} + (2\mu + 1)x\frac{\partial}{\partial x}$ $+ (\mu + 1)(t + v)\frac{\partial}{\partial t}.$
e^T	e^u	4	$\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_3 (c(T) = e^T),$ $\mathbf{W}_5 = \frac{\partial}{\partial T} + 2X\frac{\partial}{\partial X} + U\frac{\partial}{\partial U}.$	$\mathbf{W}_1, \mathbf{W}_2, \mathbf{W}_3 (c(u) = e^u),$ $\mathbf{W}_5 = \frac{\partial}{\partial u} + 2x\frac{\partial}{\partial x} + (t + v)\frac{\partial}{\partial t}.$
T^{-2}	u^{-2}	5	$\mathbf{W}_1, \mathbf{W}_2,$ $\mathbf{W}_3 (c(T) = T^{-2}), \mathbf{W}_4 (\mu = -2),$ $\mathbf{W}_6 = U^2\frac{\partial}{\partial U} + TU\frac{\partial}{\partial T} - \frac{U}{T^3}\frac{\partial}{\partial X}.$	$\mathbf{W}_1, \mathbf{W}_2,$ $\mathbf{W}_3 (c(u) = u^{-2}), \mathbf{W}_4 (\mu = -2),$ $\mathbf{W}_6 = (t + v)^2\frac{\partial}{\partial t} + u(t + v)\frac{\partial}{\partial u} - \frac{t+v}{u^3}\frac{\partial}{\partial x}.$
$T^{-\frac{2}{3}}$	$u^{-\frac{2}{3}}$	5	$\mathbf{W}_1, \mathbf{W}_2,$ $\mathbf{W}_3 (c(T) = T^{-\frac{2}{3}}), \mathbf{W}_4 (\mu = -\frac{2}{3}),$ $\mathbf{W}_7 = (XT - 3T^{\frac{2}{3}})\frac{\partial}{\partial T}$ $+ (XT^{-\frac{1}{3}} - \frac{X^2}{3})\frac{\partial}{\partial X}.$	$\mathbf{W}_1, \mathbf{W}_2,$ $\mathbf{W}_3 (c(u) = u^{-\frac{2}{3}}), \mathbf{W}_4 (\mu = -\frac{2}{3}),$ $\mathbf{W}_7 = (xu - 3u^{\frac{2}{3}})\frac{\partial}{\partial u} + (xu^{-\frac{1}{3}} - \frac{x^2}{3})\frac{\partial}{\partial x}.$

In order to determine whether a symmetry $\mathbf{W}$ of the PDE (4.24) yields a nonlocal symmetry of the nonlinear wave equation (4.20), one needs to trace back using the PDE system (4.23). Since one excludes the dependent variable V from the potential system (4.23), one needs to investigate how the variable V changes under the action induced by $\mathbf{W}$.

Note 5.2. Since $\Psi_*^{-1}(\frac{\partial}{\partial V}) = \frac{\partial}{\partial v} - \frac{\partial}{\partial t}$, where Ψ^{-1} is the inverse of the transformation Ψ (4.22), the infinitesimal components for the variables x and u keep invariant when tracing back.

Proposition 5.2. The symmetries $\mathbf{W}_6$ and $\mathbf{W}_7$ yield nonlocal symmetries of the PDE system (4.21).

Proof. Suppose the symmetry $\mathbf{W}_6$ yielded a local symmetry $\hat{\mathbf{W}}_6$ of the PDE system (4.21). Then, in evolutionary form, $\hat{\mathbf{W}}_6 = (U^2 - TUU_T + \frac{U}{T^3}U_X)\frac{\partial}{\partial U} + F([U], [V])\frac{\partial}{\partial V}$, where the function $F([U], [V])$ must depend on X, T, U, V and their derivatives. By similar reasoning as in the proof of Proposition 5.1, one can show that $F([U], [V])$ is of the form $F(X, T, U, V, U_X, U_T)$. Applying $\hat{\mathbf{W}}_6^{(\infty)}$ to the PDE system (4.21) and making appropriate substitutions, one can show that the resulting determining equation system is inconsistent. Hence, $\mathbf{W}_6$ yields a nonlocal symmetry of the PDE system (4.21).

By similar reasoning, one can show that $\mathbf{W}_7$ also yields a nonlocal symmetry of the PDE system (4.21). $\square$

Comparing Tables 5.5 and 5.7, one immediately sees that, when $c(u)$ is arbitrary, the symmetry $\mathbf{W}_3$ is a point symmetry of the PDE system (4.21), whose infinitesimal component for the variable t has an essential dependence on v . By projection, $\mathbf{W}_3$ yields a nonlocal symmetry of the nonlinear wave equation (4.20).

When $c(u) = u^{-2}$, the infinitesimal components for the variables (x, u) of the symmetry $\mathbf{W}_6$ depend on the variable v . By Note 5.2, $\mathbf{W}_6$ yields a nonlocal symmetry of the nonlinear wave equation (4.20).

When $c(u) = u^{-\frac{2}{3}}$, if the symmetry $\mathbf{W}_7$ yielded a local symmetry $\hat{\mathbf{W}}_7$ of the nonlinear wave equation (4.20), then $\hat{\mathbf{W}}_7 = \mathbf{W}_7 + f([u])\frac{\partial}{\partial t}$, where the function $f([u])$ depends on x , t , u and its derivatives. Since $\Psi_*^{-1}(\frac{\partial}{\partial v}) = \frac{\partial}{\partial v} - \frac{\partial}{\partial t}$, when tracing back to the PDE system (4.21), the infinitesimal component for the variable v must be equal to $-f([u])$. Thus $\mathbf{W}_7$ would also yield a local symmetry of the PDE system (4.21), which is a contradiction since $\mathbf{W}_7$ yields a nonlocal symmetry of the PDE system (4.21). Hence, $\mathbf{W}_7$ yields a nonlocal symmetry of the nonlinear wave equation (4.20).

Note 5.3. At a first glance, it would appear that the symmetries $\mathbf{W}_4$ and $\mathbf{W}_5$ yield nonlocal symmetries of the nonlinear wave equation (4.20). However, this is not true since the symmetries $\mathbf{W}_4$ and $\mathbf{W}_5$ yield point symmetries $\hat{\mathbf{W}}_4 = \mathbf{W}_4 + (\mu + 1)V\frac{\partial}{\partial v}$ and $\hat{\mathbf{W}}_5 = \mathbf{W}_5 + V\frac{\partial}{\partial v}$ of the PDE system (4.23) respectively since in terms of (x, t, u, v) coordinates, $\hat{\mathbf{W}}_4 = u\frac{\partial}{\partial u} + (2\mu + 1)x\frac{\partial}{\partial x} + (\mu + 1)t\frac{\partial}{\partial t} + (\mu + 1)v\frac{\partial}{\partial v} = (2\mu + 1)\mathbf{Y}_4 - \mathbf{Y}_5$ and $\hat{\mathbf{W}}_5 = \frac{\partial}{\partial u} + 2x\frac{\partial}{\partial x} + t\frac{\partial}{\partial t} + v\frac{\partial}{\partial v} = \mathbf{Y}_4 + \mathbf{Y}_7$. Hence, by projection, $\mathbf{W}_4$ and $\mathbf{W}_5$ yield point symmetries of the nonlinear wave equation (4.20).

Note 5.4. Proposition 5.2 implies that the symmetries $\mathbf{W}_6$ and $\mathbf{W}_7$ yield previously unknown nonlocal symmetries of the nonlinear wave equation (4.20).

6. Conclusion and open problems

In this paper, we presented a new systematic procedure to construct nonlocally related PDE systems for a given PDE system. The starting point for this method is the existence of a point symmetry of the given PDE system. Our method yielded previously unknown nonlocally related systems for nonlinear reaction-diffusion, nonlinear diffusion and nonlinear wave equations. In turn, it was shown that some of these new nonlocally related systems for nonlinear wave equations yielded previously unknown nonlocal symmetries.

This paper shows that the existence of a point symmetry is sufficient to generate a nonlocally related subsystem of a given PDE system. An unsolved converse problem naturally arises: Suppose a given PDE system generates a subsystem by cross-differentiation, then does the given PDE system have a point symmetry?

Potential systems are under-determined for a given PDE system with more than two independent variables. It is known that point symmetries of such potential systems can at most yield point symmetries of the given PDE system without additional gauge constraints relating potential variables and their derivatives [15]. In the case of two or more independent variables, the nonlocally related systems generated by the method presented in this paper involve natural gauge constraints due to their construction from curl-type conservation laws. Can such constrained nonlocally related systems yield nonlocal symmetries?

References

- [1] G. W. Bluman, A. F. Cheviakov, and S. C. Anco, *Applications of Symmetry Methods to Partial Differential Equations*, Appl. Math. Sci. No. 168 (Springer, New York, 2010).
- [2] A. F. Cheviakov, *Comput. Phys. Commun.* **172**, 48 (2007).
- [3] G. W. Bluman, S. Kumei, and G. J. Reid, *J. Math. Phys.* **29**, 806 (1988).

- [4] A. F. Cheviakov and G. W. Bluman, J. Math. Phys. **51**, 103521 (2010).
- [5] A. F. Cheviakov and G. W. Bluman, J. Math. Phys. **51**, 103522 (2010).
- [6] G. W. Bluman and A. F. Cheviakov, J. Math. Anal. Appl. **333**, 93 (2007).
- [7] P. A. Clarkson and E. L. Mansfield, Phys. D. **70**, 250 (1994).
- [8] P. J. Olver, *Applications of Lie Groups to Differential Equations*, GTM No. 107 (Springer, New York, 1986).
- [9] G. W. Bluman and S. Kumei, *Symmetries and Differential Equations*, Appl. Math. Sci. No. 81 (Springer, New York, 1989).
- [10] B. J. Cantwell, *Introduction to Symmetry Analysis*, Cambridge Texts in Applied Mathematics (Cambridge University Press, Cambridge, UK, 2002).
- [11] L. V. Ovsiannikov, Dokl. Akad. Nauk USSR **125**, 492 (1959) (in Russian).
- [12] G. W. Bluman and S. Kumei, J. Math. Phys. **21**, 1019 (1980).
- [13] W. F. Ames, R. J. Lohner and E. Adams, Internat. J. Nonlinear Mech **16**, 439 (1981).
- [14] G. W. Bluman and S. Kumei, J. Math. Phys. **28**, 307 (1987).
- [15] S. C. Anco and G. W. Bluman, J. Math. Phys. **38**, 3508 (1997).